
Full Length Research Paper

Evaluation of four botanical insecticides against Diamondback Moth, *Plutella Xylostella* L. (Lepidoptera: Plutellidae) on head cabbage in the central rift valley of Ethiopia

Feyissa Begna^{1*} and Tebkaw Damtew²

¹Adami Tulu Agricultural Research Center, Ziway, Ethiopia, P. O. Box 35, Ziway, Ethiopia.

²Debra Zeit Agricultural Research Center, Ziway, Ethiopia, P. O. Box 32, Debra Zeit, Ethiopia.

Accepted 6 July, 2017

Cabbage (*Brassica oleracea* var. *capitata*) is the second most important vegetable crop in Ethiopia. Many insect pest species belonging to 16 families have been recorded on cabbages. Therefore the objective of this study was to assess the efficacy of botanicals for managing diamondback moth (DBM) on head cabbage. The experiment was conducted using irrigation at Adami Tulu Agricultural Research Center (ATARC) during 2013 September to December. The head cabbage variety Copenhagen Market was used for this experiment. Treatments were arranged in randomized complete block design (RCBD) with four replications. For DBM management four locally available botanicals was sprayed six times at weekly intervals continuously. Highly significant differences among the treatments were observed after application of botanicals and chemical on DBM larvae and pupae. All botanical treatments reduced the number of DBM larval population and increased marketable yield. Among the botanicals, neem was the best treatment as it gave the highest yield and also maximum economic return.

Key words: Cabbage, botanicals, neem, cabbage, turmeric, lantana, DBM.

INTRODUCTION

Cabbage (*Brassica oleracea* L. var. *capitata*) is the second most important vegetable crop in Ethiopia with respect to production next to red pepper (*Capsicum* spp) (MOA, 2002). It is produced by private farmers (Lemma et al., 1994). The land occupied during 2010 main rainy season (Meher) was 4,802 ha with a production level of 43,483.43 tons (CSA, 2012).

Many insect pest species belonging to 16 families have been recorded in Ethiopia on head cabbage (Gashawbeza et al., 2009). However, only the diamondback moth (DBM) (*Plutella xylostella* L. Lepidoptera: Plutellidae), cabbage aphid (*Brevicoryne brassicae* L. Hemiptera: Aphididae), flea beetles (*Phylloterta* spp) and cabbage leaf miner (*Chromatomyia horticola* Goureaux) (Diptera: Agromyzidae) are of

economic importance (Tsedeke and Gashawbeza, 1994; Gashawbeza et al., 2009).

The diamondback moth is the dominant and most destructive insect pest of crucifer crops worldwide (Figure 1). Yield loss studies at Melkassa Agricultural Research Center (MACR) of the Ethiopian Institute of Agricultural Research (EIAR) showed that losses vary between 36.1 and 91.2% and complete crop failure is common in seasons of heavy infestations (Gashawbeza, 2006).

In Ethiopia, DBM pest status is believed to be strongly influenced by extensive level of insecticide usage and cabbage production methods. According to Gashawbeza and Ogol (2006), DBM is problematic in the Central Rift Valley areas where the crop is cultivated all the year-round using irrigation and where insecticide use is heavy.

Excessive use of insecticides has led to insecticidal resistance development, pest resurgence, residue hazards in foods and overall environmental contaminations. This has prompted the promotion of

*Corresponding author. E-mail: fbos2009@gmail.com. Tel.: (251) 04 64 4190 03.



Figure 1. The diamondback moth.

other DBM management alternatives such as microbial insecticide, insect growth regulators (IGRs) and botanicals. For example, aqueous extract of neem seed powder (50 g/l) and Bt (0.5k g/ha) were earlier recommended for use on cabbage under Ethiopian condition (Gashawbeza et al., 2009).

Botanical insecticides are not only effective against crop pests but remain safer to natural enemies (Patel et al., 2003). They have been in use for centuries by farmers in developing countries to control insect pests of both field crops and stored produce. Nicotine, rotenon and pyrethrum were popular among the botanical insecticides (Schmutterer, 1981). Some of these plant species possess one or more useful properties such as repellency, anti-feeding, fast knock down, flushing action, biodegradability, broad-spectrum of activity and ability to reduce insect resistance (Mochiah et al., 2011).

Therefore this study was conducted to determine the influence of botanical insecticides against DBM on head cabbage. The specific objective was:-

To assess the efficacy of some botanicals for managing diamondback moth on head cabbage.

MATERIALS AND METHODS

Description of the experimental sites

The experiment was conducted using irrigation at Adami Tulu Agricultural Research Center (ATARC) during 2013 from September to December. ATARC is located in the mid Rift Valley of Ethiopia about 167 km south from Addis Ababa. It lies at a latitude of 7° 9'N and longitude of 38° 7'E. It has an altitude of 1650 m.a.s.l. and it receives a

bimodal unevenly distributed average annual rainfall of 760.9 mm per annum. The long-term mean minimum and the mean maximum temperature are 12.6 and 27 °C respectively. The pH of the soil is 7.88. The soil is fine sandy loam in texture with sand, clay and silt in proportion of 34, 48 and 18% respectively (ATARC, 1998).

Experimental design and management

The head cabbage (*B. oleracea* var. *capitata*) variety, Copenhagen Market, was used for the present experiment. Seedlings were grown on raised seed bed of 10 m² and transplanted on October 7, 2013 when seedling reached third to fourth true leaf stage. Each plot had three ridges of four meter long and each ridges with one row of cabbage on each side. Ridges were spaced 60 cm apart. The spacing between plants was 30 cm. Treatments were arranged in randomized complete block design (RCBD) with four replications. Spacing between plots and blocks was 1 and 1.5 m, respectively. All data were collected only from the central four rows. The crop was irrigated twice per week for the first four weeks after transplanting and once weekly thereafter. Plots were fertilized with diammonium phosphate (DAP) and urea at the rate of 200 and 100 kg/ha, respectively. The whole amount of DAP was applied just before transplanting, while urea was applied by splitting the total amount in two. Half of the 100 kg was applied one month after transplanting and the remaining half at the beginning of head formation stage. Other field management practices like weeding, cultivation and maintenance of ridges were carried out as needed.

Table 1. Details of plant species used for the experiment.

Treatment code	Common name	Scientific name	Variety	Part used	Rate
1	Neem	<i>Azadirachta indica</i>	Local	kernel	50 g/L
2	Lantana	<i>Lantana camara</i>	Local	leaf	100 g/L
3	Chilli	<i>Capsicum annuum</i> (L.)	Marecofana	fruit	100 g/L
4	Turmeric	<i>Curcuma longa</i> (L.)	Local haldi	rhizome	50 g/L

Experimental materials

The experimental treatments were four botanicals and an untreated check (Table 1). Applications of treatments started three weeks after transplanting. Treatments were applied weekly until about fifteen days before harvest. Spray was made using manually operated knapsack sprayer of 15 l capacity using hollow cone nozzle. Botanical extracts were prepared one day before treatment application following the respective procedure described below. For comparison untreated check was included.

Chili extract preparation

A 200 g of red chili pepper was mixed in two liters of water and the solution was allowed to stand for 24 h, and then the solutions was strained and added 50 g of concentrate to a 3.75 l of water. The extract was later sprayed on the leaves.

Lantana camara leaf extraction

1 kg of young fresh lantana leaf was collected and crushed into small pieces using knife, then the chopped leaf was ground using grinder with 250 ml of water to make paste. The paste was strained through muslin cloth and kept for 24 h. At the time of application the aqueous extract was diluted in 9.75 l water.

Turmeric extraction

1 kg of turmeric rhizome was chopped and soaked overnight in 2 l of water. The next day the extract was filtered and filled up to 15 and sprayed on the field.

Neem kernel extraction

Neem kernel was collected from Dire Dawa, Eastern Ethiopia. Kernels were crushed in to fine powder using mortar and pestle, and sieved using wire mesh. The extract was made by mixing the powder with water in plastic container at the rate of 50 g powder per liter of water. After mixing, the solution was stirred carefully until all the powder was mixed completely with the water. This solution was left overnight. The following morning the

extract was filtered into the sprayer using plastic mesh for field use.

Data collected

Canopy spread

Canopy spread was measured with a ruler at the time of harvest. The spread of canopy was measured as the horizontal distance from one end of the plant to the other i.e. the two most outspread and directly opposite leaves of the plant (P. K. Baidoo, 2012).

Plant height

Plant height was measured from the soil surface to the apex of the plant using ruler at the time of harvest. The highest point reached by the plant was recorded as the height of the plant (Asare et al., 2010).

Yield

Marketable and unmarketable yield data were taken from the central four rows of each plot, by removing the outer damaged leaves and discarding heads with less than 4 cm in diameter. Yield losses were estimated by comparing the yield of treated cabbage with the untreated control (Judenko, 1973).

$$Yieldloss(\%) = \frac{(\bar{X} - \bar{Y})}{\bar{X}} * 100$$

Where x= mean seed cotton yield of treated plots

Y= mean seed cotton yield of untreated plots

Diamondback moth leaf damage

All plants and plant parts were examined for leaf damage by DBM before treatment application and at weekly interval thereafter. Diamondback moth leaf damage score on each leaf of a plant was taken based on a scale of 0 to 5 (0= no leaf damage; 1= up to 20% of the total leaf area damaged; 2 = 21 - 40% of the total leaf area damaged; 3 = 41 - 60% of the total leaf area damaged; 4 = 61-80% of the total leaf area damaged; and 5= more than 80 % leaf

Table 2. Mean leaf damage due to DBM on cabbage treated with different botanicals in six weeks period.

Treatment	Weeks					
	1	2	3	4	5	6
Control	3.37±0.12a	3.16±0.09 ab	3.50±.028a	4.0±0.00a	4.12±.25a	4.25±.25a
Turmeric	3.12±0.12a	2.95±0.37ab	2.10±0.4b-d	3.50±0.00ab	2.50±.28c	3.00±.41cd
Chilli	3.10±0.19a	2.91± 0.33ab	2.27±0.14b-d	2.60±0.12dc	3.25±.25b	3.25±.25bc
Lantana	3.01±0.19a	2.18±0.27ab	2.50±0.21a-d	2.70±0.75a-c	2.00±.00c	2.00±.00e
Neem	3.00±0.00a	2.50± 0.14ab	2.25±0.39b-d	2.40±0.16a-c	2.25±.25cd	2.25±.25d-e

Means followed by the same letter within a column are not significantly different (tukey's) at P = 0.05.

area damaged) (Iman et al., 1990).

Estimation of Diamondback moth population

The number of DBM larvae and pupae were recorded before and after 24 h application of botanical extracts or chemicals at weekly interval thereafter. Totally ten plants were selected randomly and examined for the presence of the different life stage of DBM. The number of larvae and pupae from each tagged leaves was counted with the help of hand lens and mean number per plant was calculated.

Stand count

Stand count after crop establishment and at harvest was taken by counting the number of plants in each plot. Number reduction in plant stand was calculated as a difference between stand counted at establishment of seedlings and harvest.

Estimation of cabbage head formation

Cabbage head formation in each treated plot was recorded during harvesting. Total number of cabbage plants with head and without head was recorded separately.

Data analysis

Data analysis was carried out using the SAS version 9.2. To stabilize the variance count and percentage data were transformed either to logarithmic or square root scale. The mean value of the recorded data's was subjected to analysis of variance (ANOVA). If there was significant difference among the treatments, mean separation was carried out using tukey's significance difference at P 0.05.

RESULTS AND DISCUSSION

Leaf damage visual scores across weeks

Leaf damage scores over six weeks period is given in Table 2. In the first week there were non-significant differences ($P < 0.05$) among all treatments, because it was before the application of any treatments. The extent of damage caused by DBM on head cabbage was almost similar, though there were leaf damage scale variations among treatments. In the 2nd week, however, there were significant differences ($P < 0.05$) among treatments in leaf damaged score. The highest leaf damage was recorded on control cabbages, whereas the least leaf damage was recorded on neem treated cabbages. Similarly in the 3rd, 4th, 5th and 6th weeks there were significant differences ($P < 0.05$) among treatments in leaf damaged score. In all the cases the control cabbage had the highest leaf damage score whereas cabbages treated with neem had the lowest leaf damage due to DBM. Cabbages treated with botanicals had intermediate leaf damage. Level of leaf damages were positively correlated with the larvae population, except in 2nd and 3rd weeks.

The present observation is in line with finding of Nakagome and Kato (1981) who stated that all crop growth stages are subjected to severe DBM infestation, so insecticide applications are required to control DBM, especially during the peak population period. When DBM is not managed the scale of leaf damage increased in untreated cabbage, but decreased generally in treated cabbages throughout the growing season. In studies made by Freddy (2011) the leaf damage was significantly lower in fields treated with insecticides than in fields not treated with insecticides. Sakai (1984) shows all crucifers suffer depredation by this pest practically throughout the growing season. Asare et al. (2010) reported the mean leaf damage for unprotected plants were higher than those which were treated in various ways.

Table 3. Mean number of DBM Larvae per plant sprayed with botanicals and chemical in 24 h post applications.

Treatment	Week					
	1	2	3	4	5	6
Control	3.16±.48a	1.50±.28ab	2.87±.51a	1.91±.41a	8.00±.57a	8.00±.57a
Turmeric	2.33±.33ab	1.79±.00ab	1.00±.57cb	1.50±.25ab	2.25±.75b	2.50±.64b
Chili	1.75±.63bc	2.00±.00a	2.00±.4ab	0.91±.4a-c	3.90±1.2b	3.92±1.2b
Lantana	2.00±.00a-c	1.25±.47a-c	1.25±.47bc	1.20±.45a-c	2.00±.57bc	2.25±.75b
Neem	0.00±.00d	1.00±.27bc	0.25±.25c	0.29±.04bc	2.00±.00b	2.25±.25b

Means followed by the same letter within a column are not significantly different (tukey's) at P = 0.05.

Table 4. Mean number of DBM Pupae per plant sprayed with botanicals and chemical in 24 h applications.

Treatment	Week					
	1	2	3	4	5	6
Control	2.75±.14a	1.62±0.21ab	1.70±.29a	1.83±0.16a	2.50±.28a	5.00±1.58a
Turmeric	1.33±.19ab	1.00±0.00 b-d	1.30±.33a	1.30±0.00a-c	1.50±.28bc	1.50±.28b
Chili	1.25±.14a-c	1.00±0.00b-d	1.00±.4ab	1.00±0.00a-d	1.00±.00c	1.0±.00b
Lantana	1.50±.00ab	0.25±0.14de	1.30±.44a	1.33±0.66a-c	1.00±.40c	0.50±.28b
Neem	1.25±.14a-b	0.50±0.28c-e	0.00±.00b	0.25±0.00cd	0.75±.25a-c	0.50±.28b

Means followed by the same letter within a column are not significantly different (tukey's) at P = 0.05.

DBM larval population 24 h after treatment application

Across all the weeks significant differences ($P < 0.05$) were observed on population of DBM larvae per plant among treatments following foliar applications (Table 3). The highest number of DBM larvae per plant were recorded from control cabbages, except during second week when the highest number of larvae per plant was recorded from chilli treated cabbages. Whereas the least number of DBM larvae were recorded from head cabbage treated with neem. Although there was reduction of DBM larval population in all treated plots 24 h after applications, the degree of DBM larval population reduction was not as expected, which might be partly attributed to the difference in pre spray larval density and to the shortest evaluation time. Within the same time span, however, the effectiveness of other botanicals was relatively variable.

In all weekly application, neem significantly reduced DBM larvae population; this was followed by lantana and turmeric. Magallona (1985) also reported that insecticides are generally considered the most effective means of protecting crops against insect damage as they provide rapid control of wide pest complex of major crucifer's pests, and growers concerned about leaf damage, even of a few holes, tend to spray insecticides. Nakagome and Kato (1981) believed that repeated insecticide applications are required to control DBM, especially during the peak population period. However, Motoyama et al. (1990) warned that effective insecticidal control of DBM might not be achieved for longer period as the insect can develop resistance to a new insecticide very

quickly because of its unique feature of insecticide resistance.

In this study, botanicals gave acceptable level of DBM larvae reduction. Nayem and Rokib (2013) found vigorous okra growth by treating with garlic bulb extracts, but not so effective than the neem extracts to control DBM. Shivanand et al. (2009) reported botanical insecticides as effective against *P. xylostella*. These plant extracts are applicable to cabbage pest management through reduction in use of synthetic insecticides spray as an important component of integrated pest management (IPM) programme. Botanical insecticides can influence the behavior and development of the herbivorous insect, which uses the plant for their reproduction as they have antifeedent, non-neuro toxic modes of action, and low environmental persistence (Arason et al., 1992). Gaby (1988) also indicated that botanicals like neem extracts play an important role in altering the attractive properties of crucifer plants to *P. xylostella*.

DBM pupae population 24 h after treatment application

Similar to the larval population, there was significant differences ($P < 0.05$) among treatments across weeks in number of DBM pupae per plant after foliar applications (Table 4). The pupal population intensity followed more or less the larval population intensity. Thus, the highest number of DBM pupa per plant was recorded from control cabbages. The least number of DBM pupae were recorded from head cabbage treated with neem. In the 2nd week there were less than one DBM pupa per plant

Table 5. Effect of botanicals on agronomic characteristics of cabbage at Adami Tullu.

Treatment	Canopy spread(cm)	Plant height(cm)	% cabbage with head	Plant stand count(number)
Control	46.68±1.56a	18.50±0.89c	83.14f	42.75±1.43b
Chilli	47.65±1.77a	19.85±0.37bc	88.47d	46.25±3.70ab
Turmeric	48.18±1.11a	19.91±0.37bc	88.75d	46.0±1.58ab
Lantana	48.05±1.20a	20.05±0.79a-c	88.42d	46.25±0.94ab
Neem	48.86±2.32a	21.77±0.64ab	91.70a	49.00±2.67ab
CV	6.6	8.01	2.75	8.6

Means followed by the same letter within a column are not significantly different (tukey's) at $P = 0.05$.

on lantana and neem treated cabbages. Similarly, in the third and fourth weeks the least number of DBM pupae were recorded from head cabbage treated with neem, turmeric and lantana treated cabbages. In the fifth week, relatively more number of pupae was recorded on all botanical treated cabbages. The numbers of DBM pupae might not be reduced across the weeks, because it is likely that more pupae would survive if there were more number of pupae in a particular treatment cabbage before treatment application. Botanicals can have effect on developmental stages of exposed pupae, which can produce morphological abnormalities in different developmental stages. Phytochemicals have considerable capacity to reduce adult emergence at low dosage, which reduce the recruitment over time and the desired characteristic of botanical insecticides. The adult emergence is affected by phytochemicals, which often cause acute and chronic toxicity in pupal stages, dead larvae-pupal intermediate stage having the head of pupa and the abdomen of a larva. Dead adults with folded wings in pupal exuvium and emerged adults were unable to escape the pupal exoskeleton, half ecdysed adults etc. (Facknath and Kawol, 1996). According to Lidet (2007) plots treated with Neem 50, Dipel and Xen Tari chemicals showed the least DBM number throughout the sampling weeks. Also Gashawbeza (2006) observed low number of DBM ranging from zero to 4 per plant in an insecticide control trial. He reported significant differences in DBM number between the untreated plot and plots treated weekly throughout the growing period.

Effect of botanicals on some agronomic characteristics

Plant height at harvest

There was significant difference ($P < 0.05$) among treatments in affecting plant height (Table 5). Cabbage sprayed with either neem produced the tallest plants. Medium plant height was measured from cabbages treated with other botanicals. However, control cabbage had the shortest plants height. This is consistent with the finding of Asare et al. (2010) who indicated that treating cabbage with insecticide reduced the insect population on cabbage and hence better growth of the crop. Nayem

and Rokib (2013) also reported that okra grows vigorously when treated with botanical insecticides.

Cabbage with heads

Significant differences ($p < 0.05$) were observed among treatments in the percentage of plants that formed head (Table 5). Cabbages treated with neem, lantana and turmeric, in decreasing order respectively formed greater percentage of heads than those cabbages treated with other botanicals. The least number of plants with head was recorded from untreated (control) plots. The DBM feed mostly on young part of the plant which is the major part for head formation. As plant losses this part they fail to form head or die under severe infestation.

Paul et al. (2001) reported that destruction of the main buds of seedlings by DBM larvae may result in plants with multiple undersized heads. Moreover, according to Asare et al. (2010) heavy head per plant was recorded for cabbages that received treatments against DBM attack when compared with the control.

Plant canopy spread

There were non-significant differences ($p > 0.05$) among treatments in plant canopy spread (table 5). Even though statistically non-significant, cabbages treated with turmeric, neem and lantana had larger diameter than cabbages treated with chilli. Moreover, although statistically non-significant plant canopy spread was negatively correlated with leaf damage and DBM larvae population except in 3rd and 4th week owing to less number of larvae recorded during those weeks. DBM larvae adversely affected the formation of head by destroying the tip of the head cabbage (Talekar and Shelton, 1993).

Effect of botanicals on cabbage yield and yield components

Effect on marketable and unmarketable yield

There were significant differences ($P < 0.05$) among treatments in marketable yield of cabbages (Table 6). Marketable yield of cabbage ranged from 27 to 92

Table 6. Effect of botanical application on yield of cabbage and economic return.

Treatment	Marketable ton/ha	Unmarketable ton/ha	Yield loss (%)	Farm gate Price birr	Gross return birr/kg	Variable cost birr/ha	Net benefit birr/ha	Marginal return rate
Control	27.75c	20.0a	-	3	83250	21096	62,154	3.92
Chilli	58.50b	8.75bc	52.52c	3	175500	28776	146,724	6.09
Turmeric	63.50b	9.50bc	56b	3	190500	25296	165,204	7.52
Lantana	63.25b	8.25bc	56b	3	189750	22776	166,974	8.31
Neem	77.25a	5.50bc	64a	3	231750	30276	201,474	7.65
CV (%)	13.57	47.61	32					

Note: Means followed by the same letter within a column are not significantly different at $p = 0.05$; Yield loss is computed as the difference between treated and untreated plots.

tons/ha. The highest level of marketable cabbage yield was obtained from plots sprayed with neem. Moreover, cabbages treated with chilli, turmeric and lantana gave comparable yield with the above mentioned botanicals. The untreated plot (control) had the lowest marketable yields. This indicates that controlling DBM populations with botanicals can double the yield of head cabbage production, even though botanicals were not equally as effective as the chemical insecticide in reducing DBM larval population and reducing associated losses.

Hasheela et al. (2010) reported that as compared to unsprayed cabbage, highest number of marketable cabbage heads was obtained from sprayed cabbage while the highest number of unmarketable cabbage heads was noted on unsprayed one. DBM larvae feeds on the marketable portions of the crop, therefore, synthetic insecticides will remain essential for the management of this pest (Hill and Foster, 2000). The plant extracts were highly effective in controlling DBM as compared to control. This could be due to the pungent smell given out by the soaked plant extract which deter animals from eating the plant (Sivapragasam and Aziz, 1990).

Yield loss

There were significant differences ($P < 0.05$) among treatments in reducing yield losses caused by DBM in cabbages (Table 6). The amount of marketable cabbage yield loss ranged from 53 to 70%. Gauging the effectiveness of control measures is one of the purposes of estimating yield losses due to pests. Thus the lowest level of yield loss relative to the control was obtained from cabbages sprayed with neem. Moreover, on chilli, turmeric and lantana treated cabbage the yield losses ranged from 52.5, 56 and 58% respectively.

Yield loss studies carried out at Melkassa Research Center of the Ethiopian Institute of Agricultural Research (EIAR) for two seasons between November 2001 and June 2002 showed that losses can vary between 36.1 and 91.2%, which corresponds to 12 and 48.7 tons/ha

(Gashawbeza 006). Complete crop failure is reported to be common on farmers' field in seasons of heavy infestation in the Central Rift Valley areas when there is no DBM management. Similarly Lidet (2007) reported that yield losses ranged between 62.8 and 74.7% which equates to 44.8 to 52.9 tons per ha at Melkassa and Wonji, respectively.

Economic return

Results of the economic analysis are presented in (Table 6). Spraying cabbage with neem gave the highest net benefit per hectare with the highest marginal return rate, but the marginal return rate of from these treatments was less than the return from lantana treatment because the application and preparation costs of lantana were less than the cost of all botanicals. Untreated plot (control) resulted in the lowest economic return with lowest marginal return rate. The economic evaluation indicated that controlling DBM population using botanicals increased net benefit and marginal return rate at least twice when compared to untreated check.

Summary and Conclusion

Leaf damage was non-significant for pre-application in 1st week. In the 2nd week, however, there were significant differences ($P < 0.05$). The highest leaf damage score was recorded on control, whereas the least leaf damage was recorded on neem treated plots. Similarly the leaf damage of the various treatments was significantly differences on 3 to 6 weeks of observations. Control cabbage, the extent of leaf damage increased across the growing season. The leaf damage on botanical treated cabbages was intermediate.

Across the weeks there were significant differences ($P < 0.05$) among treatments in affecting population of DBM larvae following foliar applications. The highest number of DBMs larvae (7 per plant) was recorded from control plots, except 2nd week in which the highest number of

larvae per plant was recorded from chilli treated plots. On the other hand, the least number of DBM larvae were recorded from head cabbage treated with neem. Within the same time span, however, the effectiveness of other botanicals was relatively variable. This shows botanical insecticide can reduce the number of DBM larvae, even though application of neem effectively controlled DBM larvae. Similar to the larval population, across weeks there were significant differences ($P < 0.05$) among treatments in number of DBM pupae per plant after foliar applications. The highest number DBM of pupa (5 per plant) was recorded from control plots, except at the 2nd week in which the highest number DBM pupae per plant were recorded from chilli sprayed plots. The least number of DBM pupae were recorded from head cabbage treated with neem. Both botanical and chemical insecticides minimized pupal population of DBM. In most case all agronomic characters, marketable yields and cabbage with head were negatively correlated with cabbage leaf damages across the week.

Significant differences were observed among treatments in some agronomic characteristics of head cabbage. Cabbages sprayed with neem produced the tallest plants and cabbages treated with other botanicals had medium plant height. However, unsprayed (control) cabbages had the shortest plants height. There were also differences among treatments in plant stand count and plants with head per plot. Large number of plant and plants with head were recorded on neem sprayed cabbages, while the least number of plant stands and plants with head per plot was observed from untreated (control) plots.

On the yield data significant differences ($P < 0.05$) among treatments was observed in marketable yield of the cabbages. The highest levels of cabbage marketable yield per plot were obtained from plots sprayed with neem foliar applications.

Finally, from this study the following recommendations have been developed

i.) To boost head cabbage production in the Central Rift Valley area, DBM and aphid that occur concurrently on head cabbage should be controlled by using neem as alternative to the currently used insecticides especially lambda cyhalothrin.

ii.) Botanical insecticide can be used to manage the population of DBM, however further studying the dose, extraction procedure, and mode of action is required.

iii.) If botanicals are used to manage DBM, they must be integrated with cabbage aphid control methods.

iv.) Botanical preparation, identification and collection are not well known by the producers in the Central Rift Valley area, so training is important for those producers.

REFERENCES

Arnanson JT, Mackinnon S, Isman MB. and urst T (1992). Insecticides

- in triopical plants with non-neurotoxic modes of action. J. of Recent Advances in Phytochem., 28. 107-108.
- Asare E, addo A and Mohammed A, 2010. Control of Diamondback Moth (*Plutella Xylostella*) On Cabbage (*Brassica Oleracea Var Capitata*) Using Intercropping with Non-Host Crops. Am. J. of Food Technol., 5(4). 269-272.
- ATARC (Adami Tulu Agricultural Research Center). 1998. ATARC Profile. Oromiya Agricultural Research Institute. Addis Ababa, Ethiopia.
- CSA (Central Statistical Agency). 2012. Report on Area and Production of Major Crops. Statistical Bulletin. Addis Ababa, Ethiopia.
- Facknath S, Kawol D (1996). Antifeedant and insecticidal effects of some plant extracts on *Crocidolomia binotalis*. J. of Insect Science and its Application 14 (5/6) 571-74.
- Freddy MO (2011). The role of predators, parasitoid and insecticides. Doctoral thesis. Uppsala, Sweden.
- Gaby S (1988). Natural crop protection based on local farm resource in the tropics and sub-tropics. Margraf Publisher Scientific Books, Germany
- Gashawbeza A (2006). Comparison of yield loss on cabbage from diamondback moth, *Plutella xylostella* (L.) (Lepidoptera: Plutellidae) using two insecticides. Crop Protection Society of Ethiopia, 25:915-919.
- Gashawbeza A, Ogol CKPO (2006). Occurrence of the Diamondback Moth (*Plutella xylostella* L.) and it's Parasitoids in Ethiopia: Influence of Geographical Region and Agronomic Traits. J. of Appl. Entomol., P 343-347.
- Gashawbeze Ayalew, Bayeh Mulatu, Mulugeta Negeri, Yeshitila Merene, Lidet Sitotaw, Ahmed Ibrahim and Tadele Tefera. 2009. Review of research on insect and mite pests of vegetable crops in Ethiopia. pp. 47-67. Cited in: Abraham Tadesse (ed.). Increasing Crop Production through Improved Plant Protection-Vol. II. Plant Protection Society of Ethiopia (PPSE). PPSE and EIAR, Addis Ababa, Ethiopia.
- Hasheela EBS, Nderitu JH, Olubayo FM, Kasina M (2010). Evaluation of cabbage varietal resistance to diamondback moth (*Plutella xylostella*) infestation and damage. Tunisian J. of Plant Protection 5: 95.
- Iman M, Dani S, Jestmandt I, Adiputra M, Ishak M (1990). Effect of insecticides on Various Field Strains of Diamondback Moth and its Parasitoid in Indonesia. P 315 In: N.S Talekar(Ed.). Diamondback Moth and Other Crucifer Pests: Proceeding of the Second International Workshop, Tainan, Taiwan, 10-14 December 1992, Asian Vegetable Research and Development Center, Shanhua, Taiwan.
- Judenko E (1973). Analytical methods for assessing yield losses caused by pests on cereal crops with and without pesticide. Tropical Bulletin No.2. UK. P. 31.
- Lemma D, Hearth E, Temesgen B, Lemaga B, Seyfu G (1994). Horticultural Research Past, Present and Future Trends. P 19-23. In: E., Hearth and Lema Dessalegne (Eds.). Proceedings of the Second National Horticultural Workshop, 1-3 December 1992, Addis Ababa, Ethiopia, IAR/FAO.
- Lidet S (2007). Distribution and management of Diamondback moth *plutella xylostella* L. (lepedoptera: plutellidae) on cabbage in the central Rift Valley of Ethiopia. An MSc Thesis Presented to the School of Graduate Studies of Haramaya University. P. 46.
- Magallona ED (1985). Development in Diamondback Moth Management in the Philippines. P 3-15. Cited in: Lidet Sitotaw, 2006. Distribution and Management of Dimandback Moth of *Pultella xylostella* L. (Lepidoptera: Plutellidae) on Cabbage in the Central Rift Valley of Ethiopia, M.Sc. Thesis, Haramaya University.
- MOA (Ministry of Agriculture)(2002). Ministry of Agriculture. Agricultural Production Statistics for the Year 2002.
- Mochiah MB, Banful B, Fening K, Amoabeng B, OffeiBonsu K, Ekyem S, Braimah H, Owusu-Akyaw M (2011). Botanicals for the Management of Insect Pests in Organic Vegetable Production. J. of Entomol. and Nematol., 3: 85-97, Available Online at [Http://Www.Academicjournals.Org/Jen](http://Www.Academicjournals.Org/Jen).
- Motoyama N, Suganuma T, Maekoshi Y (1990). Biochemical and physical characteristics of insecticide resistance in diamondback moth. P 411-418 in: N.S. Talekar (ed.). Diamondback Moth and other

- crucifer pests: Proceeding of the Second International Workshop, 11-15 December 1992, Asian Vegetable Research and Development Center, Shanhua, Taiwan.
- Nakagome T, Kato K (1981). Control of insects in cruciferous vegetables in Aichi Prefecture with special reference to diamondback moth. (In Japanese). P 79-92. In *Insects in Cruciferous Vegetables and their Control with Special Reference to Diamondback Moth*. Takeda Chemical Industries Ltd., Tokyo.
- Nayem Z, Rokib H (2013). Effects of Manually Processed Bio-pesticides on Crop Production and Pest Managements in Okra (*Abelmoschus Esculentus* (L.)). *J. of Biopesticides* 3: 7-8.
- Patel PS, Shukla NP, Patel GM (2003). Enhancing Insecticidal Properties of Cow Urine against Sucking Pests of Cotton. In: *Proceedings of the National Symposium on Front erier Areas of Entomological Research*, 5-7 November, 2003, 385.
- Paul VM, Lesley MJB, Helen C, Janny V (2001). *Discovery Learning Manual For Cabbage Pest Management*. CAB international.
- Sakai M (1984). *Chemical Control of Diamondback Moth in Japan with Special Reference to Cartap*. Agricultural Chemicals Division, Takeda Chemical Industries Ltd., Tokyo, Japan. 305p.
- Sivapragasam A, Abdul AAM (1990). Cabbage Worm on crucifers in Malaysia Cited in N. S Talekar (1990). Diamondback moth and other crucifer pest: *Proceedings of the Second International Workshop Taina, Taiwan 10-14 December, Asian Vegetable Research and Development Center*. AVRDC Publication No. 92-68 P. 75-79.
- Talekar NS, Shelton AM (1993). *Biology, Ecology and Management of Diamondback Moth*. *Ann. Rev. of Entomol.*, 38: 275-301.
- Tsedeke A, Gashawbeza A (1994). *Progress in Vegetable Management Research: 1985-1992*. P 187-189. In: E. Hearth and Lemma, D. (Eds.). *Proceedings of the Second National Horticultural Workshop*. 1-3 December 1992, Addis Ababa, Ethiopia, IAR/FAO.